

Purdue University
Purdue e-Pubs

LARS Symposia

Laboratory for Applications of Remote Sensing

1-1-1980

Forest Stand Delineation from Unsupervised Classification of Optimal Landsat Spectral, Landsat Texture and Topographic Channels

Thomas L. Logan

Alan H. Strahler

Follow this and additional works at: http://docs.lib.purdue.edu/lars_symp

Logan, Thomas L. and Strahler, Alan H., "Forest Stand Delineation from Unsupervised Classification of Optimal Landsat Spectral, Landsat Texture and Topographic Channels" (1980). *LARS Symposia*. Paper 387.
http://docs.lib.purdue.edu/lars_symp/387

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Reprinted from

Symposium on

Machine Processing of

Remotely Sensed Data

and

Soil Information Systems

and

Remote Sensing and Soil Survey

June 3-6, 1980

Proceedings

The Laboratory for Applications of Remote Sensing

Purdue University
West Lafayette
Indiana 47907 USA

IEEE Catalog No.
80CH1533-9 MPRSD

Copyright © 1980 IEEE
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org.

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

FOREST STAND DELINEATION FROM UNSUPERVISED CLASSIFICATION OF OPTIMAL LANDSAT SPECTRAL, LANDSAT TEXTURE AND TOPOGRAPHIC CHANNELS

THOMAS L. LOGAN

Jet Propulsion Laboratory/California
Institute of Technology

ALAN H. STRAHLER

University of California

Landsat data, in conjunction with collateral data sources such as synthesized texture channels and digital terrain models, can be used to delineate forest stands and other ecological land units found in the coniferous North American forest environment. Incorporation of texture and terrain channels enhances site-specific stratification of Landsat data, promoting delineation of forest stand units of a size and homogeneity approaching those found on manually prepared maps used in the management of timber, range, wildlife, watershed and recreational resources.

The procedure is a joint research effort between the Jet Propulsion Laboratory of the California Institute of Technology and the University of California at Santa Barbara. The classification approach includes 1) compressing Landsat spectral data into one or two new channels of data using ratio and principle components techniques; 2) generating two texture measures where one channel emphasizes tonal contrast derived from statistical texture techniques and the other emphasizes spatial extent and shape using image segmentation procedures; 3) processing National Cartographic Information Center - U.S. Geological Survey Digital Terrain information into elevation, slope and aspect channels; 4) reducing the number of synthesized channels by using divergence analysis to identify channels not contributing significantly to the separation of preliminary training classes; 5) introducing spatial constraints by including line and sample coordinates into the unsupervised classification algorithm; and 6) properly weighting selected spectral, texture and terrain channels such that no single data set overpowers the others in unsupervised classification.

The combination of spectral tone, tonal texture, spatial texture, topographic data and line and sample location

coordinates, is likely to be sufficient for the stand delineation task because each contributes a separate, independent piece of information towards the stand delineation problem. Spectral tone is most important for recognizing the existence of a feature and combines with the topographic data to provide species information. Tonal texture measures the neighborhood contrast of spectral tones providing an indication of relative timber volume. Spatial texture stratifies tone to quantify the spatial extent and shape of tonal patterns. The topographic information provides a powerful independent parameter well known to improve forest classification accuracies because of its ecological predictive effect. Inclusion of line and sample coordinates introduces a strong spatial constraint designed to permit analyst regulation over the automatic merging of distant and unrelated, but similar appearing features.

Target area for generation of maps delineating forest stands and related ecological land units is the 220 square kilometer Doggett Creek watershed located in the Klamath National Forest of northern California. The mountainous topography ranges from 500 to 2100 meters in elevation, and bears a variety of important coniferous timber types including douglas fir, ponderosa pine, white and red fir, and several miscellaneous hardwoods such as black oak and madrone.

Part of the research described in this paper was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under NASA Contract NAS 7-100.

CH1533-9/80/0000-0338 \$00.75 © 1980 IEEE

1980 Machine Processing of Remotely Sensed Data Symposium